



## The Reproduction Properties of *Serranus cabrilla* (L., 1758) and *Serranus hepatus* (L., 1758) from Izmir Bay

Dilay Birim, Bulent Cihangir\*, E. Mumtaz Tirasin

Dokuz Eylül University, Institute of Marine Sciences and Technology. 35340 Inciralti-Izmir

+905515913101

bulent.cihangir@deu.edu.tr

---

### Abstract

In this study, reproduction and some biological aspects of *Serranus cabrilla* (comber) and *Serranus hepatus* (brown comber) from Izmir Bay were investigated. The fish material was collected between February 2007 and September 2008 during demersal trawl surveys carried out by R/V K. Piri Reis which is operated by Institute of Marine Sciences and Technology of DEU. Totally, 554 females and 340 hermaphrodite *S. cabrilla* specimens, and 443 females and 2070 hermaphrodite *S. hepatus* specimens were examined. The observed reproduction periods and first maturity lengths for *S. cabrilla* were between March-May and 129 mm fork length, and for *S. hepatus* March-September and 77 mm total length. The mean batch fecundity were found 2869 and 587 eggs for *S. cabrilla* and *S. hepatus*, respectively. Fork lengths of *S. cabrilla* ranged between 95-210 mm and total lengths of *S. hepatus* ranged between 63-119 mm. Both species showed negative allometric growth. Length-weight relationship changed in sampling periods and for gender for both species.

### Keywords:

*Serranus cabrilla*, *Serranus hepatus*, reproduction, hermaphroditism

---