



## Morphometric Differences Between *Serranus cabrilla* (Linnaeus, 1758) Populations from Mediterranean and Aegean Sea

Deniz Yaglioglu\*<sup>1,2</sup>, Deniz Ayas<sup>3</sup>, Ali Uyan<sup>4</sup>, Servet Dogdu<sup>4</sup>

<sup>1</sup>Department of Biology, Faculty of Arts and Science, Düzce University, Düzce, Turkey

<sup>2</sup>Biodiversity Implementation and Research Center (DU-BIYOM), Düzce University, Düzce, Turkey

<sup>3</sup>Department of Seafood and Processing Technology, Faculty of Fisheries, Mersin University, Mersin, Turkey

<sup>4</sup>Iskenderun Technical University, Marine Science and Technology Faculty, Iskenderun, Hatay, Turkey.

\*denizyaglioglu@duzce.edu.tr,

Phone: +90.536.6417798

### Abstract

The aim here is to investigate morphometric structure of comber *Serranus cabrilla* populations from Aegean Sea (Izmir) and Mediterranean Sea (Mersin). Truss measurements were made on the specimens by collecting X-Y co-ordinate data for 10 morphological landmarks. Discriminant function analysis was used to investigate morphometric differences between populations using SPSS. Univariate statistics (ANOVA) revealed highly significant differences among locations from 9 out of 18 truss measurements. In discriminant function analysis, a high degree of morphologic differentiation was detected among populations. Proportions of correctly classified individuals into their original grouping showed high classifications in the Mediterranean Sea and Aegean Sea samples, 100 % and 97.1%, respectively. It was observed that samples from the Aegean Sea (Izmir) and Mediterranean Sea (Mersin) were different from each other.

### Keywords:

Serranidae, Comber, *Serranus cabrilla*, morphometric

---