



## **-RESEARCH ARTICLE-**

### **Assessing DNA Barcodes for Identification of Pufferfish Species (Tetraodontidae) in Turkish Marine Waters**

Cemal Turan<sup>\*</sup>, Mevlüt Gürlek, Deniz Ergüden, Ali Uyan, Serpil Karan, Servet A. Dođdu

Molecular Ecology and Fisheries Genetic Laboratory, Marine Sciences and Technology Faculty, Iskenderun Technical University, 31220, Iskenderun, Hatay, Turkey.

#### **Abstract**

In Turkish marine waters, pufferfish belongs to Tetraodontidae family are represented with 8 species, *Lagocephalus lagocephalus*, *L. sceleratus*, *L. spadiceus*, *L. suezensis*, *L. guentheri*, *Spherooides pachygaster*, *Torquigener flavimaculosus* and *Tylerius spinosissimus*. DNA barcoding can be useful in the assessment of cryptic or morphologically similar species of identification which is widespread in marine environment. The DNA barcode identification of the eight puffer species of the Tetraodontidae family in Turkish marine waters were examined by using mtDNA sequencing of the amplified partial mitochondrial cytochrome c oxidase I (COI) gene. COI contained 189 variable and 337 conservative nucleotides of which 183 were parsimony informative over 526 bp. Mean genetic diversity of all species was found to be 0.18164. The highest (0.26127) and lowest (0.00305) nucleotide divergence was observed between *L.spadiceus* and *T. flavimaculosus* and between *L. spadiceus* and *L. guentheri*, respectively. The number of different haplotypes were 12 out of 23 sequences, and there was no shared haplotypes between pufferfish species.

#### **Keywords:**

Pufferfish, molecular identification, DNA Barcoding, COI

#### **Article history:**

Received 17 December 2017, Accepted 19 December 2017, Available online 19 December 2017

---

<sup>\*</sup> Corresponding Author: Cemal Turan, e-mail: cemal.turan@iste.edu.tr

## Introduction

Pufferfishes are marine fish species that are distributed in tropical and subtropical areas of the Atlantic, Indian and Pacific Ocean. Puffers include 28 genera and approximately 184 species in all over the world marine waters within the Tetraodontidae family (Matsuura, 2015; Farrag et al., 2016), among which at least ten are found in the eastern Mediterranean (Farrag, 2014). This indigenous invasive species has established large populations along the coasts of many countries of the eastern Mediterranean basin such as Israel, Lebanon, Turkey (Mediterranean and Aegean coasts), Cyprus and Greece (Aegean and Ionian coasts), while still rapidly expanding westwards along the coasts of Egypt, Libya, and along the entire Tunisian coastline (Soussi et al. 2014). Apart from several large species used for human consumption as a delicious food in few countries, particularly in China, Korea, Japan and Taiwan (Oyaizu et al. 2000), most pufferfish species have not commercial value. Besides the small size of most species, the family is renowned for the occurrence of a powerful toxin in their skin and organs called tetrodotoxin (TTX). Tetrodotoxin is a very potent neurotoxin and one of the strongest marine paralytic toxins (El-Sayed et al., 2003; Sato et al., 2008).

In Turkish marine waters, pufferfishes are represented with 8 species, *Lagocephalus lagocephalus* (Linnaeus, 1758), *Lagocephalus sceleratus* (Gmelin, 1789), *Lagocephalus spadiceus* (Richardson, 1845), *Lagocephalus suzensis* Clark & Gohar, 1953, *Lagocephalus guentheri* Miranda Ribeiro, 1915, *Sphoeroides pachygaster* (Müller & Troschel, 1848), *Torquigener flavimaculosus* Hardy & Randall, 1983, *Tylerius spinosissimus* (Regan, 1908) (Turan et al., 2007).

Molecular genetic studies on mtDNA have proven benefits for examining the phylogeny and phylogeography of marine species (Meyer, 1993; Avise, 1994; Turan et al. 2015a). Sequence analysis of mtDNA regions is a quick tool to reveal phylogenetic relationships of marine species (Avise, 1994; Turan et al. 2008; Tabata & Taniguchi, 2000). Ever since different regions of mtDNA evolve at different rates, specific mtDNA regions have been targeted for inter and intra specific variation (Hauser et al. 2001; Mohindra et al., 2007; Turan et al., 2015b). DNA barcoding is a global venture that provides a standardized and effective genetic marker to marine and freshwater biodiversity, with significant conservation applications. The DNA barcoding approach is concentrated on a single part of the mitochondrial genome, because it presents portions conserved across taxa that are appropriate for primer design, while including polymorphism between and within species (Hebert et al., 2003; Kress & Erickson, 2008). The cytochrome oxidase subunit I (COI) region of the mitochondrial genome is sufficiently diverse so as to let the specific identification of a great majority of fish species (Kochzius et al., 2008; Kochzius et al., 2010).

Simple identification of pufferfishes by DNA barcoding and current level of interspecific and intraspecific genetic variation at pufferfish species which distributed in Turkish waters are very important to know. In spite of the wide scientific interest given to this family because of their ecological impact and having tetrodotoxin in their tissue, there have been no remarkable study which investigated genetic identification and structure of these species in Turkish waters.

The goal of this study is to evaluate the practicability of DNA barcoding in the monitoring species biodiversity distributed along the Turkish marine waters at two levels by confirming the taxonomic identification and specifying intraspecific and interspecific variations for eight pufferfish species found in Turkish marine waters.

## Material and Methods

Species, *Lagocephalus lagocephalus*, *L. sceleratus*, *L. spadiceus*, *L. suezensis*, *L. guentheri* and *Torquigener flavimaculosus*, were collected from the Antalya and Iskenderun Bay. The picture of pufferfish species collected from the Antalya and Iskenderun Bay are shown at Figure.1. All the samples were put in plastic bags individually and frozen at -20 °C till they were transported to the laboratory. All tissue samples were stored at -20 °C and 95 % ethanol till the analysis.




Figure 1. Pufferfish species in Turkish Marine Waters.

Total genomic DNA was extracted from the muscle and fin samples using the DNeasy Blood and Tissue Kit (Qiagen, USA). Manufacturer's protocols were used during all steps. Polymerase chain reaction (PCR) amplification was performed with following selective primers especially designed for this experiment:

COI-Forward 5'-TCAACCAACCACAAAGACATTGGCAC-3'

COI-Reserved 5'-ACTTCAGGGTGACCGAAGAATCAGAA-3'

The PCRs were conducted in a 50 ml total volume with 0.4 uM of each primer, 0.2 mM of dNTP and 1.25U of Taq DNA polymerase in a PCR buffer that included 20mM of Tris-HCl (pH



|              |          |          |          |          |          |          |          |          |           |
|--------------|----------|----------|----------|----------|----------|----------|----------|----------|-----------|
| <b>Hap10</b> | -        | -        | -        | -        | -        | -        | -        | <b>1</b> | <b>1</b>  |
| <b>Hap11</b> | -        | -        | -        | -        | -        | -        | -        | <b>1</b> | <b>1</b>  |
| <b>Hap12</b> | -        | -        | -        | -        | -        | <b>3</b> | -        | -        | <b>3</b>  |
| <b>Total</b> | <b>2</b> | <b>5</b> | <b>5</b> | <b>5</b> | <b>2</b> | <b>3</b> | <b>2</b> | <b>3</b> | <b>27</b> |




Figure 2. Minimum spanning tree that shows the relationships among the haplotypes.

Table 2. Variable nucleotide positions of COI DNA barcode in pufferfish species. The DNA barcode variable nucleotides are indicated, while identity is shown by dashes.

|        | 10  | 20 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | 100 |
|--------|---|----|----|----|----|----|----|----|----|-----|
|        | *   | *  | *  | *  | *  | *  | *  | *  | *  | *   |
| Hap_1  | CCCCACAATAATGTCTATACAGACAAGTGATCGAGCCACATCCTCCCTTTGCTTCCACTCGAGCTAAGCCTCCAGCCACTATACCTCGCTCGTTACCACC  |    |    |    |    |    |    |    |    |     |
| Hap_2  | TTA.G...G.C.CTC.C.TGAT..TAC.G.T.CA..GT.C.TC.TA.CCT...T.G.G.A..T.GCAT..G...TC..CA..T...C.A.AT....      |    |    |    |    |    |    |    |    |     |
| Hap_3  | TTA.G...G.C.CTC.C.TGAT..TAC.G.T.CA..GT.C.TC.TA.CCT...T.G.G.A..T.GCAT..G...TC..CA..T...C.A.AT....      |    |    |    |    |    |    |    |    |     |
| Hap_4  | .....C...C.G...TTGT.CA.C.....CC.....TT...A....CT..CA.CA..G.C...TAT...GG...T.                          |    |    |    |    |    |    |    |    |     |
| Hap_5  | GTA.....C..TC.C...AT..TACA...ACA.....C.TA.CCT...TAGTG.A..T.GCTT.CG...TC...G..A..ATC.ACGT.A...         |    |    |    |    |    |    |    |    |     |
| Hap_6  | GTA.G...G.C.CTC.C.TGAT..TAC.G.T.CA..GT.C.TC.TA.CCT...T.G.G.A..T.GCAT..G...TC..CA..T...C.A.AT....      |    |    |    |    |    |    |    |    |     |
| Hap_7  | ...T..TG...CACA..C.T..TTTTA..CC..CA...T...TC.TAC.CC..AGTG.GGTGTTT.CT..CAT.AA.T.C.CCTAC.C.CTACATA.C.T  |    |    |    |    |    |    |    |    |     |
| Hap_8  | ...T..TG...CACA..C.T..TTTTA..CC..CA...T...TC.TAC.CC..AGTG.GGTGTTT.CT.GCAT.AA.T.C.CCTAC.C.CTACATA.C.T  |    |    |    |    |    |    |    |    |     |
| Hap_9  | .T.T.T..G.CA..C.C.T..TTTTA..CC..CAT...G.TGC...CGC...AAA.TG.TG.A..CC..CG.CAA.T...GTTTC.C.C.T.ATT.T...  |    |    |    |    |    |    |    |    |     |
| Hap_10 | .T.T.T..G.CA..C.C.T..TTTTA..CC..CAT...G.TGC...CGC...AAA.TG.TG.A..CC..CG.CAA.T...GTTTC.C.C.T.AT..T...  |    |    |    |    |    |    |    |    |     |
| Hap_11 | .T.T.T..G.CA..C.C.T.ATTTA..CC..CAT...G.TGC...CGC...AAA.TG.TG.A..CC..CG.CAA.T...GTTTC.C.C.T.AT..T...   |    |    |    |    |    |    |    |    |     |
| Hap_12 | ...T.T..C.CC..TC...T.ATATTACACC.A.A.TG..CT.CT.T..CATG..A.TATTG.T..C.T...TGCATGAC.GT.ACTA..TT.GT.GG... |    |    |    |    |    |    |    |    |     |

Continued

|  | 110  | 120 | 130 | 140 | 150 | 160 | 170 | 180 |
|--|--|-----|-----|-----|-----|-----|-----|-----|
|  | *  | *   | *   | *   | *   | *   | *   | *   |
|  | TTCCCCATCCCCGACTACTTAAACACTCTCGACACGCTCTCTCCCCCGCGCCGCCACTCCACAACCCGAGAGCGCTCACCACCTGTT  |     |     |     |     |     |     |     |
|  | CG.T.T.C.A.TA.TCCTC.C.G....GT.G.C.....T.....TTAGC...T...CTT.TTT...AC...TAAC....G.....    |     |     |     |     |     |     |     |
|  | CG.T.T.C.A.TA.TCCTC.C.G....GT.G.C.....T.....TTAGC...T.A..CTTTTTT...AC...TAAC.T..G.....   |     |     |     |     |     |     |     |
|  | AG.T.T.C..T..G...T.....AG.G.A..TCT.TTTGT.AT.T..TT..TA.....A..G.TA...T.....               |     |     |     |     |     |     |     |
|  | CG...T.C.ATTA.TCCTT..C..T...GT.GTTTATCAAT.GAA.ATAGC.A.T...TCTT.TTT...AC.G.TAAC.....C..   |     |     |     |     |     |     |     |
|  | CG.T.T.C.A.TA.TCCTC.C.G....GT.G.C.....T.....TTAGC...T...CTT.TTT...AC...TAAC....G.....    |     |     |     |     |     |     |     |
|  | AGG...T.AATTA.AA.TC.C...GA...A.TC.AT.TA..G..TATAA.TA..A.C.A..T..T.TT..AGAGA.CA..G.AGGTCC |     |     |     |     |     |     |     |
|  | AGG...T.AATTA.AA.T.CCG..GA...A.TC.AT.TA..G..TATAA.TA..A.C.A..T..T.TT..AGAGA.CA....GGTCC  |     |     |     |     |     |     |     |
|  | CGAT...AA...AG.TC....GAA.GT...GTAT.TCT.GT..A.ATC.TAT..C.C..CT.C.T.A..G..ATCT.T....A..    |     |     |     |     |     |     |     |
|  | CGAT...AA...AG.TC....AA.GT...GTAT.TCT.GT..A.ATC.TAT..C...CT.T.T.A..G...TCT.T....A..      |     |     |     |     |     |     |     |
|  | CGAT...AA...AG.T.....AA.GT...GTAT.TCT.GT..A.ATC.TAT..C...CT.T.T.A..G...TCT.T....A..      |     |     |     |     |     |     |     |
|  | GG..TT..TATT.GT..TC.G..T.A.TATA.GCTAT..CTCT.A.ATAG.TA.T...A..C.CCT.T.T...T.AC..T..T..A.. |     |     |     |     |     |     |     |

Species special DNA barcode were detected whereas common DNA barcode was not detected for all species. Kimura 2 parameter method (Kimura, 1980) was selected as a best method for intra and interspecific variations. Mean genetic diversity for all species was found to be 0.18164. The matrix of pairwise distances within species is presented in Table 3. The intraspecific genetic diversity within *L. sceleratus*, *L. suezensis*, *L. lagocephalus*, *L. guentheri* and *S. pachygaster* was observed to be zero while it was highest within *T. flavimucolus* (0.01149). The lowest genetic distance is observed between *L. guentheri* and *L.spadiceus* (0.00305) whereas the highest one is observed between *T. flavimucolus* and *L.spadiceus* (0.26127).

Table 3. The matrix of intraspecific genetic distances between species (below diagonal) and genetic diversity (transversal diagonal given in bold).

|                             | 1              | 2              | 3              | 4              | 5              | 6              | 7              | 8              |
|-----------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <i>L. sceleratus</i> (1)    | <b>0.00000</b> |                |                |                |                |                |                |                |
| <i>L.spadiceus</i> (2)      | 0.19850        | <b>0.00229</b> |                |                |                |                |                |                |
| <i>L. suezensis</i> (3)     | 0.13133        | 0.22827        | <b>0.00000</b> |                |                |                |                |                |
| <i>L. lagocephalus</i> (4)  | 0.21867        | 0.10202        | 0.19204        | <b>0.00000</b> |                |                |                |                |
| <i>L. guentheri</i> (5)     | 0.19673        | 0.00305        | 0.22639        | 0.09859        | <b>0.00000</b> |                |                |                |
| <i>T. flavimucolus</i> (6)  | 0.25622        | 0.26127        | 0.24380        | 0.24518        | 0.26042        | <b>0.01149</b> |                |                |
| <i>T. spinosissimus</i> (7) | 0.21345        | 0.21469        | 0.20354        | 0.20343        | 0.21338        | 0.19274        | <b>0.00896</b> |                |
| <i>S. pachygaster</i> (8)   | 0.25101        | 0.24912        | 0.23208        | 0.21013        | 0.24759        | 0.25409        | 0.21512        | <b>0.00000</b> |

The Neighbour Joining and Maximum Parsimony phylogenetic approaches resulted in similar tree topologies. In Neighbour joining phylogenetic tree, two main phylogenetic nodes were detected; in the first main node, *T. flavimucolus* and *T. spinosissimus* grouped together. In second main node, 3 branches were detected; *S. pachygaster* was in the first branch, *L. spadiceus*, *L. guentheri* and *L. lagocephalus* were grouped in the second branch on which *L. guentheri* and *L. spadiceus* were grouped together as a sister group, and *L. sceleratus* and *L. suezensis* were grouped as a third branch (Figure 3).






Figure 3. Neighbour joining phylogenetic tree based on COI sequences. Fish drawings from Froese & Pauly (2016).

In the Maximum Parsimony phylogenetic tree, two phylogenetic nodes were detected; in the first node, *T. flavimucolus* and *T. spinosissimus* were grouped together. In second node 3 branches were detected; *S. pachygaster* was in the first branch, *L. spadiceus*, *L. guentheri* and *L. lagocephalus* were grouped as a second branch in which *L. guentheri* and *L. spadiceus* were grouped together as a sister group, and *L. sceleratus* and *L. suezensis* were grouped as a third branch (Figure 4).






Figure 4. Maximum Parsimony phylogenetic tree based on COI sequences. Fish drawings from Froese & Pauly (2016).

### Discussion

In the present study, the DNA barcoding of the eight pufferfish species which are distributed in the Turkish marine waters were investigated. All the species under the three genera were clearly separated in the NJ and MP trees with a high bootstrap value. The universal primers successfully amplified the target region in all species, generating 27 COI barcodes of 526 bp. Common haplotypes was not detected between species, and the DNA barcode sequences clearly discriminated taxonomic status of all pufferfish species examined.

Genetic diversity within species were calculated as zero for *L. sceleratus*, *L. suezensis*, *L. lagocephalus*, *L. guentheri* and *S. pachygaster*. This low genetic diversity may be explained low number of samples sequenced in the present study. On the other hand, the detected low genetic diversity may be due to founder effect that is usually observed in the alien species invation which

form established populations starting from meager individuals. A similar result is reported by Keskin & Atar (2013) using DNA barcoding to identify 89 commercially important freshwater and marine fish species found in Turkish ichthyofauna. Vinas & Tudela (2009) studied genetic identification of eight Scombrid species using mtDNA control region, mtDNA COI gene and nuclear DNA ITS1 region, and reported that credibility of COI gene is questionable that also reported that COI gene is not a good marker for inferring evolutionary relationships in Thunnus species.

The present study support the power of COI for species identification which is in accordance with other similar studies. Lakra et al. (2011) investigated DNA barcoding of fish and marine life representing 79 Genera and 37 Families from the Indian Ocean using cytochrome c oxidase I gene (COI) of the mtDNA and concluded that morphological characters were strongly authenticated the efficacy of COI in identifying the fish species with designated DNA barcodes that make DNA barcoding approach successful. Kochzius et al. (2010) aimed to evaluate the applicability of the three mitochondrial genes 16S rRNA (16S), Cytochrome b (Cyt b), and cytochrome oxidase subunit I (COI) for the identification of 50 European marine fish species by combining techniques of DNA barcoding and microarrays. As a result, while Cyt b and COI are equally well suited for DNA barcoding of fishes. On the other hand, 16S has drawbacks in discriminating closely related species. All these studies have shown that genetic identification by COI barcodes can provide a useful tool to identify species and to detect possibly cryptic species, and even to describe new species.

In conclusion, in this study has strongly authenticated the efficacy of COI in identifying the pufferfish species with designated barcodes. The present results also suggest that COI barcoding can be used as pragmatic method for resolving unambiguous identification of pufferfish species in marine waters of Turkey with applications in its management and conservation.

### Acknowledgement

Thanks to Republic of Turkey Ministry of Food, Agriculture and Livestock General Directorate of Agricultural Research and Policies (TAGEM-16/AR-GE/21) for financial support.

### References

- Avise, J.C. 1994. Molecular Markers, Natural History and Evolution. Chapman and Hall, New York, USA, 511 pp.
- El-Sayed, M., Yacout, G. A., El-Samra, M., Ali, A., & Kotb, S. M. 2003. Toxicity of the Red Sea pufferfish *Pleuranacanthus sceleratus* "El-Karad". *Ecotoxicology and environmental safety*, 56(3), 367-372.
- Farrag, M. M. S. 2014. Fisheries and Biological studies on Lessepsian pufferfish, *Lagocephalus sceleratus* (Gmelin, 1789) (Family: Tetraodontidae) in the Egyptian Mediterranean waters. Faculty of Sciences Al-Azhar University (Assiut), Egypt.
- Farrag, M., El-Haweet, A. A., and Moustafa, M. A. 2016. Occurrence of puffer fishes (Tetraodontidae) in the eastern Mediterranean, Egyptian coast-filling in the gap. *BioInvasions Record*, 5(1).
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, 39(4), 783-791.
- Froese, R., & Pauly, D. 2016. FishBase. Worldwide web electronic publication. 2014.

- Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41: 95-98.
- Hauser, L., Turan, C., Carvalho, G.R. 2001. Haplotype frequency distribution and discriminatory power of two mtDNA fragments in a marine pelagic teleost (Atlantic herring, *Clupea harengus*). *Heredity*, 87: 621–630.
- Hebert, P.D.N., Cywinska, A., Ball, S.L., deWaard, J.R. 2003. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B* 270: 313–321.
- Keskin, E., & Atar, H. H. 2013. DNA barcoding commercially important fish species of Turkey. *Molecular Ecology Resources*, 13(5), 788-797.
- Kimura, M. (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16: 111-120.
- Kochzius, M., Nölte, M., Weber, H., Silkenbeumer, N., Hjörleifsdottir, S., Hreggvidsson, G. O., Marteinson, V., Kappel, K., Planes, S., Tinti, F., Magoulas, A., Garcia Vazquez, E., Turan, C., Hervet, C., Campo Falgueras, D., Antoniou, A., Landi, M., Blohm, D. 2008. DNA microarrays for identifying fishes. *Marine Biotechnology*, 10: 207–217.
- Kochzius, M., Seidel, C., Antoniou, A., Botla, S. K., Campo, D., Cariani, A., Vazquez, E. G., Hauschild, J., Hervet, C., Hjörleifsdottir, S., Hreggvidsson, G., Kappel, K., Landi, M., Magoulas, A., Marteinson, V., Nölte, M., Planes, S., Tinti, F., Turan, C., Venugopal, M.N., Weber, H. Blohm, D. 2010. Identifying fishes through DNA barcodes and microarrays. *PLoS One*, 5 (9): 1-15.
- Kress, W.J., Erickson, D.L. 2008. DNA barcodes: Genes, genomics, and bioinformatics. *PNAS*: 105 (8): 2761–2762.
- Kumar, S., Stecher, G., & Tamura, K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular biology and evolution*, 33(7), 1870-1874.
- Lakra, W. S., Verma, M. S., Goswami, M. Lal, K. K., Mohindra, V., Punia, P., Gopalakrishnan, A., Singh, K. V. R., Ward, D., & Hebert, P. 2011. DNA barcoding Indian marine fishes. *Molecular Ecology Resources*, 11: 60–71.
- Landi, M., Dimech, M., Arculeo, M., Biondo, G., Martins, R., Carneiro, M., Carvalho, G.R., Brutto, S.L., & Costa, F. O. 2014. DNA barcoding for species assignment: the case of Mediterranean marine fishes. *PLoS One*, 9(9), 1-9.
- Matsuura, K. 2015. Taxonomy and systematics of tetraodontiform fishes: a review focusing primarily on progress in the period from 1980 to 2014. *Ichthyological Research*, 62(1), 72-113.
- Meyer, A. 1993. Evolution of mitochondrial DNA in fishes. In: *Biochemistry and Molecular Biology of Fishes*. Elsevier Science Publishers, 2: 1-38.
- Mohindra, V., Singh, R.K., Palanichamy, M., Ponniah, A.G., Lal, K.K. 2007. Genetic identification of three species of the genus *Clarias* using allozyme and mitochondrial DNA markers. *Journal of Applied Ichthyology*, 23:104–109.
- Nei, M., & Kumar, S. 2000. *Molecular evolution and phylogenetics*. Oxford university press.
- Oyaizu, M., Fujimoto, Y., Takenaga, F., & Itoh, S. 2000. Fatty acid composition of total lipids in puffer fish meat. *Food Preservation Science*, 26(6), 333-338.
- Sanger, F., Nicklen, S., Coulson, A.R. 1977. DNA Sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences*, 74:5463-5467.
- Santini, F., Nguyen, M. T. T., Sorenson, L., Waltzek, T. B., Lynch Alfaro, J. W., Eastman, J. M., & Alfaro, M. E. 2013. Do habitat shifts drive diversification in teleost fishes? An example

- from the pufferfishes (Tetraodontidae). *Journal of Evolutionary Biology*, 26(5), 1003-1018.
- Sato, K., Akai, S., Shoji, H., Sugita, N., Yoshida, S., Nagai, Y., Suzuki, K., Nakamura, Y., Kajihara, Y., Funabashi, M., & Yoshimura, J. 2008. Stereoselective and efficient total synthesis of optically active tetrodotoxin from d-glucose. *The Journal of Organic Chemistry*, 73, 1234–1242.
- Souissi, J. B., Rifi, M., Ghanem, R., Ghazzi, L., Boughedir, W., & Azzurro, E. 2014. *Lagocephalus sceleratus* (Gmelin, 1789) expands through the African coasts towards the Western Mediterranean Sea: a call for awareness. *Management*, 5(4), 357-362.
- Steinke, D., Connell, A. D., & Hebert, P. D. 2016. Linking adults and immatures of South African marine fishes. *Genome*, 59(11), 959-967.
- Tabata, K., Taniguchi, N. 2000. Differences between *Pagrus major* and *Pagrus auratus* through mainly mtDNA control region analysis. *Fisheries Science*, 66: 9-18.
- Thompson, J. D., Higgins, D.G., Gibson, T.J. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22: 4673–4680.
- Turan, C., Ergüden, D., Çevik, C., Gürlek, M., Turan, F. 2015a. Molecular systematic analysis of shad species (*Alosa* spp.) from Turkish marine waters using mtDNA genes. *Turkish Journal of Fisheries and Aquatic Sciences*, 15 (1): 149-155.
- Turan, C., Erguden, D., Gürlek, M., Yaglioglu, D., & Keskin, Ç. 2007. Atlas and systematics of marine Bony fishes of Turkey. Nobel, Adana, Turkey.
- Turan, C., Gunduz, I, Gurlek, M., Yaglioglu, D. 2008. Systematics of Scorpaenidae species in the Mediterranean Sea inferred from mitochondrial 16S rDNA sequence and morphological data. *Folia Biologica*, 57: 219-226.
- Turan, C., Gurlek, M., Erguden, D., Yaglioglu, D., Ozturk, B., Uyan, A., Reyhaniye, A. N., Ozbalcilar, B., Erdogan, Z. A., Ivanova, P., Soldo, A. 2015b. Population Genetic Analysis of Atlantic Bonito *Sarda sarda* (Bloch, 1793) using Sequence Analysis of mtDNA D-Loop Region. *Fresenius Environmental Bulletin*, 45 (3): 231-237.
- Viñas, J., & Tudela, S. 2009. A validated methodology for genetic identification of tuna species (genus *Thunnus*). *PLOS one*, 4(10), 1-10.
- Wang, Z. D., Guo, Y. S., Liu, X. M., Fan, Y. B., & Liu, C. W. 2012. DNA barcoding South China Sea fishes. *Mitochondrial DNA*, 23(5), 405-410.