



**-RESEARCH ARTICLE-**

**Morphological Anomalies Observed on *Vertebralina striata* d'Orbigny, 1826 Test in the Northern Coast of Karaburun Peninsula (Izmir-Turkey)**

Engin Meriç<sup>1</sup>, Niyazi Avşar<sup>2</sup>, Mehmet Baki Yokeş<sup>3\*</sup>, Feyza Dinçer<sup>4</sup>

<sup>1</sup> Moda Hüseyin Bey Sokak No: 15/4 34710 Kadıköy-İstanbul/Turkey

<sup>2</sup>Çukurova Üniversitesi, Mühendislik-Mimarlık Fakültesi, Jeoloji Mühendisliği Bölümü, 01330 Balcalı-Adana/Turkey

<sup>3</sup>AMBRD Doğa Bilimleri, Hanımefendi Sokak, No: 160/6, 34384 Şişli-İstanbul/Turkey

<sup>4</sup>Nevşehir University, Mühendislik ve Mimarlık Fakültesi, Jeoloji Mühendisliği Bölümü, 50300 Nevşehir/Turkey

**Abstract**

Morphological abnormalities have been observed on the test of recent foraminifer samples and locally different ecological conditions, such as, hypersalinity, presence of heavy metals and thermal springs are suggested to be the cause of this phenomenon. Abnormal test development, as well as twins and triplets are commonly observed in natural and artificial salt pans which form hypersaline environments. Submarine freshwater springs can cause extreme seasonal salinity variations on the coastal regions which affects the benthic foraminifer species and results in the abnormal test development. *Vertebralina striata* d'Orbigny individuals with abnormal aperture morphology have been found in samples collected around the thermal springs in Çeşme-Ilıca Bay. More surprisingly, specimens with similar aperture abnormality have also been observed in samples collected from other two stations on the Karaburun Peninsula. Five of the abnormal specimens had two apertures the others had three. A specimen collected from Ilıca Bay had two apertures, one of which is the typical *Vertebralina striata* d'Orbigny aperture, but the other was distantly located and rounded. *Vertebralina striata* d'Orbigny is widely distributed in world's seas, however such aperture abnormalities has not been reported elsewhere. Observation of similar morphological abnormalities in three distinct locations suggests a common cause. The submarine spring found in Ilıca Bay supports the idea that spring waters affect the chemical composition of the seawater locally, leading to this aperture abnormality.

**Keywords:**

*Vertebralina striata*, foraminifera, morphological abnormality, Karaburun, Turkey.

\* Corresponding Author: Mehmet Baki Yokeş, e-mail: bakiyokes@gmail.com

**Article history:**

Received 01 February 2019, Accepted 13 May 2019, Available online 16 May 2019

---

**Introduction**

Hypersalinity, presence of heavy metals, thermal springs, chemicals carried by the coastal and submarine springs found in karstic regions and rapid change in ecological parameters are suggested to play a major role in changing the ecological conditions that leads to abnormal test morphology in foraminifers (Boltovskoy et al., 1991). Abnormal test development, as well as twins and triplets are commonly observed in natural and artificial salt pans which form hypersaline environments (Almogi-Labin et al., 1992; Stouff et al., 1999; Geslin et al., 2000; Debenay et al., 2001). Submarine freshwater springs can cause extreme seasonal salinity variations on the coastal regions which affect the benthic foraminifer species and result in the abnormal test development (Nigam et al., 2006; Meriç et al., 2008a). The high heavy metal concentrations, either natural or artificial, locally affect the benthic foraminifera. The heavy metals directly disposed from the industrial plants or carried by the water systems result in the change in environmental conditions and cause variations on test morphology in benthic foraminifers (Alve, 1991; Sharifi et al., 1991; Yanko et al., 1998 and 1999; Meriç et al., 2002b and 2005), as well as abnormal test development (Geslin et al., 2002; Meriç et al., 2003b; Elberling et al., 2003; Meriç et al., 2008e and 2009a).

The benthic foraminifer assemblages in thermal springs were investigated at shallow habitats in the Aegean Sea (Thierman et al., 1997; Varnavas et al., 1999). Similar studies were conducted around the thermal springs in Ilica Bay - Çeşme (Izmir, western coast of Turkey), (Meriç, 1986; Avşar & Meriç, 2001). *Peneroplis* individuals with abnormal morphology are highly abundant around the submarine springs (Sözeri, 1966; Sellier de Civrieux, 1970).

*Vertebralina striata* d'Orbigny individuals with abnormal aperture morphology have been found in samples collected around the thermal springs in Ilica Bay. More surprisingly, specimens with similar aperture abnormality have also been observed in samples collected from another station on west side of Karaburun Peninsula. Such aperture abnormality has not been reported elsewhere. Observation of similar morphological abnormalities in two distinct locations could suggest a common cause.

**Material and Methods**

Sediment samples were manually collected on 06.11.2008, around a submarine spring which is located in Çeşme-Ilica Bay (Figure 1, Station 1), on the northwestern of Karaburun Peninsula (coordinates: 4240949N and 0444185E). The spring was taken as the center and three transects were set on A (210°N), B (120°N) and C (290°N) directions (depth range 1-4 m). Sampling was done at each 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90 and 100 m distances according to the spring. But, because of a jetty, samples beyond 40 m could not be obtained on line A.




Figure 1. Map of the study area and locations of the stations. Station 1, Çeşme- Ilıca Bay; Station 2, west side of Karaburun Peninsula; Station 3, northeast side of Karaburun Peninsula.

45 surface sediment samples were collected from west of Karaburun Peninsula on 07.11.2018, at a depth of 1.00-19.30 m (Figure 1, Station 2). Starting from the center (coordinates: 4261268N, 0448465E), three transects were set in A (210°), B (125°) and C (290°) directions. Sampling was done at each 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90 and 100 m distances according to the center.

45 surface sediment samples were also collected from northeast of Karaburun Peninsula on 07.11.2018, at a depth of 1.00-5.70 m (Figure 1, Station 3). Starting from the center (coordinates of the center: 4276724N, 0458535E), three transects were set in A (80°), B (10°) and C (160°) directions. Sampling was done at each 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90 and 100 m. distances according to the center.

For each sample five grams of wet sediment were weighed and treated with 10 % H<sub>2</sub>O<sub>2</sub> for 24 hours. The samples were washed with pressurized water on 0.063 sieve and dried at 50°C oven. The dried samples were further sieved with 2.00, 1.00, 0.500, 0.250, 0.125 mm mesh sizes. The foraminifer individuals were separated under binocular microscope.

SEM microphotographs were taken in T.P.A.O. Research Center (Ankara) with Jeol. JSM-6490 LV microphotography instrument.

## SYSTEMATICS

Superfamily Nubecularioidea Jones, 1875

Fischerinidae Millet, 1898

Subfamily Nodobaculariellinae Bogdanovich, 1981

Genus *Vertebralina* d'Orbigny, 1826

(Figures 2 and 3)

Description of *V. striata* (from Loeblich and Tappan, 1988)

Test is flattened, broad, and somewhat elongated, early chambers are slightly trochospiral and involute, final chamber is uncoiled and rectilinear; wall is calcareous, imperforate, porcellaneous, with numerous closely spaced longitudinal costae on the surface; aperture is terminal, lip is a narrow elongate slit with smooth and thickened bordering, which is slightly turned toward the side with the umbilical view of the coil. The species is known from Holocene with a distribution range in Atlantic, Pacific and Mediterranean.

**Remarks**

The benthic foraminifera *V. striata* has a global distribution in tropical and subtropical seas. It has been recorded from Indian Ocean (Azazi, 1992; Hottinger et al., 1993; Parker & Gischler, 2011; Langer et al., 2013; Montaggioni, 1981), from Pacific (Millet, 1898; Cushman, 1917 and 1932; Graham and Militante, 1959; McCulloch, 1977; Cheng & Zheng, 1978; Baccaert, 1987; Haig, 1988; Loeblich & Tappan, 1988; Hatta & Ujiie, 1992; Loeblich & Tappan, 1994; Yassini & Jones, 1995; Cann et al., 2000; Richardson & Clayshuttle, 2003; Debeney, 2003; Makled & Langer, 2011) and also from Atlantic (Williamson, 1858; Cushman, 1929).

In the Mediterranean, it has been recorded from Adriatic (Wiesner, 1923; Daniels, 1970; Cimerman & Langer, 1991); Gulf of Naples (Sgrrella and Moncharmont-Zei, 1993); Mallorca (Milker & Schmiedl, 2012). *V. striata* is also commonly found along the Turkish coastline, on the eastern Aegean coast, it has been frequently observed in Gulf of Saros, Dardanelles, Gökçeada, Bozcaada, Lesvos Island, Gulf of Edremit, Ayvalık, Dikili, Çandarlı, Gulf of İzmir, Karaburun Peninsula, Kuşadası, Gulf of Güllük, Gulf of Gökova, Datça and Marmaris (Meriç & Avşar, 2001; Meriç et al., 2002a and b, 2003a and b, 2004; 2008a, and b; 2009a, b and c). It has also been recorded from all along the Mediterranean coast of Turkey, from Kalkan, Kaş, Kekova, Antalya Bay to Gulf of Iskenderun (Avşar et al., 2001; Avşar et al., 2008; Meriç et al., 2008c).

**Results**

Nine specimens show typical morphological characteristics of *V. striata*, except the aperture morphology. Five of the abnormal specimens have two apertures (Figure 2 and Figure 3) and the others have three (Figure 2). Test and aperture morphologies of two normal *V. striata* specimens are shown in Figure 2, which were collected from Station 1, Line-B, distance 10 m, depth 3.8 m and Line-C, distance 5 m, depth 2.50 m, respectively. A specimen found in




Figure 2. (1) *Vertebralina striata* d'Orbigny: a and b, side views, Station 1, Line-B, distance 10, depth 3.80 m.; (2) *Vertebralina striata* d'Orbigny. Side view of the aperture, Station 1, Line-C, distance 5, depth 2.50 m.; (3) *Vertebralina striata* d'Orbigny. Side view of the abnormal test, Station 1, Line-A, distance 25, depth 3.20 m.; (4) *Vertebralina striata* d'Orbigny. Abnormal test

and aperture, a, side view, b, detailed view of two apertures, c, detailed view of second aperture. Station 1, Line-C, distance 15, depth 1.80 m.; (5) *Vertebralina striata* d'Orbigny. a, side view of the test and b, detailed view of two apertures, Station 3, Line-B, distance 80, depth 3.20 m.; (6) *Vertebralina striata* d'Orbigny. a, side view and b, detailed view of two aperture, Station 2, Line-C, distance 20, depth 2.00 m.; (7) *Vertebralina striata* d'Orbigny. a, side view, b, detailed view of two apertures, Station 2, Line-C, distance 30, depth 2.30 m. (Scale bar: 100 $\mu$ m).




Figure 3. (1) *Vertebralina striata* d'Orbigny. a, side view, b and c, detailed views of two apertures, Station 2, Line-C, distance 45, depth 2.70 m.; (2) *Vertebralina striata* d'Orbigny. Side view, Station 1, Line-B, distance 35, depth 4.10 m.; (3) *Vertebralina striata* d'Orbigny. a, side view and b, detailed view of the aperture, Station 1, Line-C, distance 5, depth 2.50 m.; (4) *Vertebralina striata* d'Orbigny. a, side view, b, detailed views of the aperture, Station 2, Line-C, distance 35, depth 2.40 m.; (5) *Vertebralina striata* d'Orbigny. a, side view, and b, detailed view of the aperture, Station 1, Line-C, distance 60, depth 1.30 m. (Scale bar: 100 $\mu$ m).

Station 1, Line-A, distance 25 m, depth 3.20 m shows abnormal chamber morphology (Figure 2), whereas, another specimen collected from Line-C, distance 15 m, depth 1.80 m has two apertures, one of which is the typical *V. striata* aperture, but the other is distantly located and rounded (Figure 2). On the other hand, *V. striata* individual found on Line-B, distance 80.00 m at a depth of 3.20 m in Station 3 has two apertures (Figure 2). Another specimen with two apertures

were found in Station 2 on Line-C, distance 20 m at a depth of 2.00 m. (Figure 2) and two more specimens with similar abnormalities were found in Station 2, on Line-C, distance 30.00 m at a depth of 2.30 m (Figure 2). Individuals with two apertures were also observed in Station 2, on Line-C, distance 45.00 m at a depth of 2.70 m (Figure 3). Individuals with three distinct apertures were found in Station 2, Line-B, distance 35.00 m at a depth of 4.10 m (Figure 3), Line-C, distance 35.00 m at a depth of 2.40 m (Figure 3) and in Station 1, Line-C, distance 5.00 m at a depth of 2.50 m and Line-C, distance 60 m at a depth of 1.30 m.

The presence of Pacific or Red Sea originated alien species, such as *Euthymonacha polita* (Chapman) (Meriç et al., 2010) and *Coscinospira acicularis* (Batsch) (Meriç et al., 2011) in certain locations in the Aegean Sea suggests the presence of different environmental conditions specific to these locations. In contrast to the abundant populations of *Amphistegina lobifera* Larsen along the coasts of Gulf of Kuşadası and both stations on Karaburun Peninsula, no individual of this species have been observed around the submarine spring in Ilıca Bay. Besides, togethernesses observed between *Peneroplis planatus* (Fichtel and Moll) and *Coscinospira hemprichii* Ehrenberg, *Peneroplis pertusus* (Forskal) and *C. hemprichii* in Ayvalık (Meriç et al., 2008d), as well as, togethernesses observed between *V. striata* and *C. hemprichii* in Gulf of Kuşadası (Meriç et al., 2009d) supports the presence of location-specific environmental conditions hypothesis.

## Discussion

Individuals with aperture abnormalities do not show any other morphological anomaly, suggesting that the factors affecting the aperture abnormality do not act on all of the test, but only on certain characteristics, such as aperture morphology.

*V. striata* is widely distributed in world seas. However, the morphological abnormalities observed in samples from Ilıca Bay and Karaburun peninsula have not been reported from elsewhere, which suggests that these abnormalities may be specific to the locality. The observation of colored tests and morphological abnormalities other than aperture morphology have been observed in samples from the same locations (Meriç et al., 2012a, 2012b), supporting the idea that the thermal springs located around (Erişen et al., 1996; Akkuş et al., 2005) may affect the chemical composition of the sea water locally, leading to test abnormalities. The samples from Ilıca Bay have been collected near a submarine spring. Although there is no known submarine springs around the stations on the Karaburun Peninsula, observation of abnormal *V. striata* test suggests their presence.

## Acknowledgement

Authors would like to thank Nihat BOZDOĞAN, Erhan YILMAZ and Physist Tuğrul TÜZÜNER (T.P.A.O. Research Center, Ankara) for their support in SEM (Jeol. JSM-6490 LV) microphotography.

## References

- Akkuş İ., Akıllı H., Ceyhan S., Dilemre A. and Tekin Z. (2005) *Türkiye Jeotermal Kaynakları Envanteri*. M. T. A. Genel Müdürlüğü, Ankara.
- Almogi-Labin A., Perelis-Grossowicz L. and Raab M. (1992) Living Ammonia from a hypersaline inland pool. Dead Sea Area, Israel. *Journal of Foraminiferal Research*, 22 (3): 257-266.
- Alve E. (1991) Benthic foraminifera in sedimen cores reflecting heavy metal pollution in Sorfijord, Western Norway. *Journal of Foraminiferal Research*, 21 (1): 1-19.



- Asano K. (1951) 14. Rotaliidae. In L.W. Stach (ed.) *Illustrated Catalogue of Japanese tertiary smlaeer foraminifera*. Japan: Tokyo, Hosokawa Printing Co. 1-21.
- Avşar N. and Meriç E. (2001) Çeşme-Ilıca Koyu (İzmir) bölgesi güncel bentik foraminiferlerinin sistematik dağılımı. *H. Ü. Yerbilimleri*, 24: 13-22.
- Avşar N., Meriç E. and Ergin M. (2001) İskenderun Körfezi bentojenik sedimentlerinin foraminifer içeriği. *H. Ü. Yerbilimleri*, 24, 97-112.
- Avşar N., Meriç E., Alramazanoğlu A. and Dinçer F. (2008) Antalya Körfezi (GB Türkiye) kıyı bölgesi bentik foraminifer toplulukları. *H. Ü. Yerbilimleri*, 29 (3), 111-136.
- Azazi G. (1992) Recent seafloor benthonic foraminiferal analysis from the Gulf of Suez, Egypt. In Takayanagi Y. and Saito T. (eds.) *Studies in Benthic Foraminifera: Proceedings of the Fourth International Symposium on Benthic Foraminifera, Sendai, 1990*. Japan: Tokai University Press, 135-149.
- Baccaert J. (1987) *Distribution patterns and taxonomy of benthic foraminifera in the Lizard Island Reef Complex, northern Great Barrier Reef, Australia*. Ph. D. Thesis, Univ. Liege: C.A.P.S. Lab. Biosedimentologie.
- Boltovskoy E., Scott D. B. and Medioli F. S. (1991) Morphological variations of benthic foraminiferal tests in response to changes in ecological parameters: A review. *Journal of Paleontology*, 65 (2): 175-185.
- Cann J. H, Belperio A. P. and Colin V. M. W. (2000) Late Quaternary paleosealevels and paleoenvironments inferred from foraminifera, northern Spencer Gulf south Australia. *Journal of Foraminiferal Research*, 30 (1), 29-53.
- Cimerman F. and Langer M. R. (1991) *Mediterranean foraminifera*. Slovenska Akademija Znanosti in Umetnosti, Akademia Scientiarum et Artium Slovenica, Ljubljana.
- Cushman J. A. (1917) New species and varieties of foraminifera from the Philippines and adjacent waters. *Proceedings of the United States National Museum*, 51: 651-662.
- Cushman J. A. (1929) The foraminifera of the Atlantic Ocean, Part VI, Miliolidae, Ophalmidiidae and Fischerinidae. Washington. *United States National Museum, Bulletin*, 104 (6), 129 p.
- Cushman J. A. (1932) The foraminifera of the tropical Pacific collections of the "Albatross", 1899-1900. Pt. 1. Astorhizidae to Trochamminidae, *Bulletin of the United States National Museum*, 161, i-vi, 1-88.
- Çakmakoğlu A. and Bilgin Z. R. (2006) Karaburun Yarımadası'nın Neojen öncesi stratigrafisi. *Maden Tetkik ve Arama Dergisi*, 132, 33-62.
- Daniels, C. H. Von. (1970) Quantitative ökologische analyse der zeitlichen und raumlichen verteilung rezenter foraminiferen im Limski-kanal bei Rovinj (nördliche Adria). *Göttinger Arbeiten für Geologie und Paläontologie*. 8, 1-109, pls. 1-8, Göttingen.
- Debenay J.-P., Geslin E., Eicher B. B., Duleba V., Sylvestre F. and Eicher P. (2001) Foraminiferal assemblages in a hypersaline Lagoon, Araruama (R.J.) Brazil. *Journal of Foraminiferal Research*, 31 (2): 133-151.
- Elberling B., Knudsen K. L., Kristansen P. H. and Asmund G. (2003) Applying foraminiferal stratigraphy as a biomarker for heavy metal contamination and Mining impact in a fiord in West Greenland. *Marine Environmental Research*, 55: 235-256.
- Erişen B., Akkuş İ., Uygur N. and Koçak A. (1996) *Türkiye Jeotermal Envanteri*. MTA Genel Müdürlüğü, Ankara.
- Geslin E., Stouff V., Debenay J.-P., and Lesourd M. (2000) Environmental variation and foraminiferal test abnormalities. In Martin R.E. (ed.) *Environmental Micropaleontology*,



- New York: Kluwer Academic/Plenum Publishers, pp. 191-215. [Topics in Geobiology vol. 15].
- Geslin E., Debenay J.-P., Duleba W. and Bonetti C. (2002) Morphological abnormalities of foraminiferal tests in Brizalian environments; comparison between polluted and nonpolluted areas. *Marine Micropaleontology*, 45: 151-168.
- Graham J. J. and Militante P. J. (1959) Recent foraminifera from the Puerto Galera area, northern Mindoro, Philippines. *Stanford University Publications, Geological Sciences*, 6 (2), 1-171.
- Haig D. W. (1988) Miliolid foraminifera from inner neritic sand and mud facies of the Papuan Lagoon, New Guinea. *Journal of Foraminiferal Research*, 18, 203-236.
- Hatta A. and Ujiie H. (1992) Benthic foraminifera from Coral Seas between Ishigaki and Iriomote Islands, Southern Ryukyu Island Arc, Northwestern Pasific. *Bulletin of the College of Science, University of the Ryukyus*, 53, 49-119; 54, 163-287.
- Hottinger L., Halicz E. and Reiss Z. (1993) *Recent foraminiferida from the Gulf of Aqaba, Red Sea*. Ljubljana: Slovenska Akademija Znanosti in Umetnosti, Academia Scientiarum et Artium Slovenica.
- Loeblich Jr. A. R. and Tappan H. (1988) *Foraminiferal genera and their classification*. New York: Van Nostrand Reinhold Company.
- Loeblich Jr. A. R. and Tappan H. (1994) *Foraminifera of the Sahul Shelf and Timor Sea*. Cushman Foundation for Foraminiferal Research, Special Publication, No: 31.
- McCulloch I. (1977) *Qualitative observations on recent foraminiferal tests with emphasis on the eastern Pacific: Parts I-III*. Los Angeles University of Southern California.
- Meriç E. (1986) Deniz dibi termal kaynakların canlı yaşamına etkisi hakkında güncel bir örnek (Ilica-Çeşme-İzmir). *Türkiye Jeoloji Kurumu Bülteni*, 29 (2): 17-21.
- Meriç E. and Avşar N. (2001) Benthic foraminiferal fauna of Gökçeada Island (North Aegean Sea) and its local variations. *Acta Adriatica*, 42 (1): 125-150.
- Meriç E., Avşar N. and Nazik A. (2002a) Bozcaada (Kuzey Ege Denizi) bentik foraminifer ve ostrakod faunası ile bu toplulukta gözlenen yerel değişimler. *Ç. Ü. Yerbilimleri (Geosound)*, 40-41: 97-119.
- Meriç E., Avşar N. and Bergin F. (2002b) Midilli Adası (Yunanistan-Kuzeydoğu Ege Denizi) bentik foraminifer faunası ve bu toplulukta gözlenen yerel değişimler. *Ç. Ü. Yerbilimleri (Geosound)*, 40-41: 177-193.
- Meriç E., Avşar N., Bergin F. and Barut İ. F. (2003a) Edremit Körfezi (Kuzey Ege Denizi, Türkiye) bentik foraminifer topluluğu ile ekolojik koşulların incelenmesi. *Ç. Ü. Yerbilimleri (Geosound)*, 43: 169-182.
- Meriç E., Avşar N., Bergin F. and Barut İ. F. (2003b) A note on three abnormal samples of benthic foraminifers from the Dikili Bay (Turkey) in northeastern Aegean Sea: *Peneroplis planatus* (Fichtel ve Moll), *Rosalina* sp. ve *Elphidium crispum* (Linné). *Bulletin of the Mineral Research and Exploration*, 127: 1-14.
- Meriç E., Avşar N. and Bergin F. (2004) *Benthic foraminifera of Eastern Aegean Sea (Turkey) Systematics and Autoecology*. Istanbul: Turkish Marine Research Foundation and Chamber of Geological Engineers of Turkey, Publication No: 18.
- Meriç E., Avşar N., Nazik A., Alpar B., Yokeş B., Barut İ. F. and Ünlü S. (2005) Gemlik Körfezi'nin yüzey çökellerinin foraminifer, ostrakod ve mollusk faunası, foraminifer kavkılarında gözlenen morfolojik anomaliler ile sedimentolojik, hidrokimyasal ve biokimyasal özellikleri. *Maden Tetkik ve Arama Dergisi*, 131: 21-48.

- Meriç E., Avşar N., Nazik A., Tunoğlu C., Yokeş B., Barut İ. F., Yücesoy-Eryılmaz F., Tuğrul B., Görmüş M., Öncel M. S., Orak H., Kam E. and Dinçer F. (2008a) Harmantaşı Mevkii (Saros Körfezi-Kuzey Ege Denizi) deniz içi kaynaklar çevresindeki foraminifer ve ostrakod topluluğuna bu alandaki çevresel koşulların etkisi. *Maden Tetkik ve Arama Dergisi*, 136: 63-84.
- Meriç E., Avşar N., Yokeş B. and Dinçer F. (2008b) Alibey ve Maden adaları (Ayvalık-Balıkesir) yakın çevresi bentik foraminiferlerinin taksonomik dağılımı. *Maden Tetkik ve Arama Dergisi*, 137. 49-65.
- Meriç E., Avşar N., Nazik A., Yokeş B. and Dinçer F. (2008c) A review of benthic foraminifers and ostracodes of the Antalya coast. *Micropaleontology*, 54 (3-4), 187-197.
- Meriç E., Yokeş M.B., Nielsen J. K., Görmüş M., Avşar N. and Dinçer F. (2008d) Abnormal formations in peneroplid foraminifers: Peneroplis-Coscinospira togetherness. *Anales de Biologia*, 30, 1-7.
- Meriç E., Avşar N., Yokeş B., Tuğrul A. B., Bayarı S., Özyurt N., Barut İ. F., Balkis N., Uysal K. and Kam E. (2008e) Morphological abnormalities in benthic foraminifers of the Antalya coast. *Micropaleontology*, 54 (3-4): 241-276.
- Meriç E., Avşar N., Mekik F., Yokeş B., Barut İ.F., Dora Ö., Suner F., Yücesoy-Eryılmaz F., Eryılmaz M., Dinçer F. and Kam E. (2009a) Alibey ve Maden Adaları (Ayvalık-Balıkesir) Çevresi Genç Çökellerinde Gözlenen Bentik Foraminifer Kavkılarındaki Anormal Oluşumlar ve Nedenleri. *Türkiye Jeoloji Bülteni*, 52(1): 31-84.
- Meriç E., Avşar N., Nazik A., Yokeş B., Ergin M., Eryılmaz M., Yücesoy-Eryılmaz F., Gökaşan E., Suner F., Tur H., Aydın Ş. and Dinçer F. (2009b) Çanakkale Boğazı'nın güncel bentik foraminifer, ostrakod, mollusk topluluğunu denetleyen faktörler ile çökel dağılımının jeokimyası. *Türkiye Jeoloji Bülteni*, 52 (2), 155-215.
- Meriç E., Avşar N., Barut İ. F., Yokeş M. B., Taş S., Eryılmaz M., Dinçer F. and Bircan C. (2009c) Opinions and comments on the benthic foraminiferal assemblages observed around the mineral submarine springs in Kuşadası (Aydın-Turkey). In Barbieri F. (ed) *13th Congress RCMNS: 2-6 September 2009, Naples, Italy*. Acta Naturalia de "L'Ateneo Parmense", 45(1/4): 222-223.
- Meriç E., Yokeş M.B., Avşar N. and Bircan C. (2009d) A new observation of abnormal development in benthic foraminifers: *Vertebralina-Coscinospira* togetherness. *Marine Biodiversity Records*, 2, e167, doi:10.1017/S1755267209990893.
- Meriç E., Yokeş M. B., Avşar N. and Bircan C. (2010) An oasis for alien benthic foraminifera in the Aegean Sea. *Aquatic Invasions*, 5 (2), 191-195.
- Meriç E., Yokeş MB and Avşar N (2011) A new guest in Ilıca Bay (Çeşme-Izmir-Turkey), *Coscinospira acicularis*. *Marine Biodiversity Records*, 4(e94):1-5, doi:10.1017/S1755267211000492.
- Meriç, E., Avşar, N., Nazik, A., Yokeş, M., Barut, İ. F., Eryılmaz, M., Kam, E., Taşkın, H., Başsarı, A., Dinçer, F., Bircan, C., Kaygun, A., 2012a, Ilıca Koyu (Çeşme-İzmir) bentik foraminifer-ostrakod toplulukları ile Pasifik Okyanusu ve Kızıldeniz kökenli göçmen foraminiferler ve anormal bireyler. *M.T.A. Dergisi*, 145, 62-78, Ankara.
- Meriç, E., Avşar, N., Nazik, A., Yokeş, Dora, Ö., Barut, İ. F., Eryılmaz, M., Dinçer, F., Kam, E., Aksu, A., Taşkın, H., Başsarı, A., Bircan, C., Kaygun, A., 2012b, Karaburun Yarımadası kuzey kıyılarının oşinografik özelliklerinin bentik foraminifer ve ostrakod toplulukları üzerindeki etkileri. *M.T.A. Dergisi*, 145, 22-47, Ankara.

- Millett F. W. (1898) Report on the recent foraminifera of the Malay Archipelago collected by Mr. A. Durrand, F. R. M. S. Part III. *Journal of the Microscopical Society*, 1898: 607-614.
- Nigam R., Saraswat R. and Kurtarkar S. R. (2006) Laboratory experiment to study the effect of salinity variations on benthic foraminiferal species-*Pararotalia nipponica* (Asano). *Journal Geological Society of India*, 67: 41-46.
- Sellier De Civrieux J. M. (1970) Mutaciones recientes del género *Peneroplis* y relaciones filogénicas con otros Soritidae. *Revista Espanola de Micropaleontologia*, 11 (1): 5-12.
- Sgarrella F. and Moncharmont-Zei M. (1993) Benthic foraminifera of the Gulf of Naples (Italy): systematics and autoecology. *Bulletino della Societa Paleontologica Italiana*. 32 (2), 145-264.
- Sharifi A. R. L., Croudace W. and Austin R. L. (1991) Benthic foraminiferids as pollution indicators in Southampton Water, southern England. United Kingdom. *Journal of Micropaleontology*, 10 (1): 109-113.
- Sözeri B. (1966) İzmir, Çeşme Ilıcası plaj kumundaki aktuel foraminiferler ve varyasyonları. *Türkiye Jeoloji Kurumu Bülteni*, 10 (1-2): 143-154.
- Stouff V., Debenay J.-P. and Lesourd M. (1999) Origin of morphological abnormalities in *Ammonia* (Foraminifera): Studies in laboratory and natural environments. *Journal of Foraminiferal Research*, 29 (2): 152-170.
- Thierman F., Akoumianaki L., Hughes J. A. and Giere O. (1997) Benthic fauna of a shallow-water gaseohydrothermal vent area in the Aegean Sea (Milos, Greece). *Marine Biology*, 128 (1): 149-159.
- Varnavas S. P., Halbach P., Halbach M., Panagiotaras D., Rahders E. and Hubner A. (1999) Characterization of hydrothermal fields and hydrothermal evolution in the Hellenic Volcanic Arc. In Balopoulos E.Th., Chronis G.Th., Lipiatou E. and Oliounine I. (eds) *International Conference on Oceanography of the Eastern Mediterranean and Black Sea, Scientific Report. Athens, Greece, 23-26 February 1999*. Luxembourg: Office for Official Publications of the European Communities, 343-344.
- Wiesner H. (1923) Die Milioliden der östlichen Adria. Prag-Bubenec. 113 p.
- Williamson V. C. (1858) *On the recent foraminifera of Great Britain*. Ray Society, London
- Yassini I. and Jones B. G. (1995) *Foraminiferida and Ostracoda from estuarine and shelf environments on the southeastern coast of Australia*. Wollongong: The University of Wollongong Press.
- Yanko V., Ahmad M. and Kaminsky M. (1998) Morphological deformities of benthic foraminiferal tests in response to pollution by heavy metals: Implications for pollution monitoring. *Journal of Foraminiferal Research*, 28 (3): 177-200.
- Yanko V., Arnold A. and Parker W. (1999) The effect of marine pollution on benthic foraminifera: In Sen Gupta, B (ed.) *Modern Foraminifera*. London: Kluwer Academic Publishers, 217-235.
- Zheng Shou-Yi, T. C., Cheng X., Vang T. and Fu X. Z. (1978) The Quaternary foraminifera of the Dayuzhang irrigation area, Shandong Province, and a preliminary attempt at an interpretation of its depositional environment. *Studia Marina Sinica*, 13, 16-78.