

Natural and Engineering Sciences

Abstracts Book of International Grouper Workshop, 07-08 October 2016, Bodrum, Turkey.

Length-Weght Relationship of Brown Comber, *Serranus hepatus*, Linnaeus, 1758 from Iskenderun Bay, Northeastern Mediterranean, Turkey

Asiye Başusta¹, Nuri Başusta¹, Levent Sangün²

¹Fisheries Faculty, Firat University, 23119 Elazig, Turkey
²Adana Vocational School , Çukurova University, 01330, Adana, Turkey
*Corresponding Author: Nuri Başusta, e-mail: nbasusta@hotmail.com, Tel: +905445184995

Abstract

This study describes the length-weight relationship (LWR) for *Serranus hepatus* species from Iskenderun Bay, Northeastern Mediterranean, Turkey. A total of 202 *S. hepatus* individuals were caught from Iskenderun Bay, during the 2015-2016 fishing season. The total length ranged from 5.7-9.5cm and body weight 2.23-12.19 g for *S. hepatus*. Length-weight relationship was found as the W= $0.0172L^{2.9664}$ (R²=0.84, F_{1,200}=21.221, P < 0.001) and SE_b= 0.091 (t-test: 32.706 p < 0.01) for all individuals of *S. hepatus*. The types of growth were found isometric growth for all individuals of *S. hepatus*.

Keywords:

Length-weight relationship, Brown comber, Serranus hepatus, northeastern Mediterranean Sea